



# Homogenization of a model of compressible miscible flow in porous media

Yacine Y. Amirat, K. Hamdache, A. Ziani

## ► To cite this version:

Yacine Y. Amirat, K. Hamdache, A. Ziani. Homogenization of a model of compressible miscible flow in porous media. RR-0991, INRIA. 1989. inria-00075568

**HAL Id: inria-00075568**

**<https://hal.inria.fr/inria-00075568>**

Submitted on 24 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



UNITÉ DE RECHERCHE  
IRIA-ROCQUENCOURT

Institut National  
de Recherche  
en Informatique  
et en Automatique

Domaine de Voluceau  
Rocquencourt  
BP 105  
78153 Le Chesnay Cedex  
France  
Tel (1) 39 63 55 11

# Rapports de Recherche

N° 991

*Programme 7*

## HOMOGENIZATION OF A MODEL OF COMPRESSIBLE MISCIBLE FLOW IN POROUS MEDIA

Youcef AMIRAT  
Kamel HAMDACHE  
Abdel Hamid ZIANI

Mars 1989



# **HOMOGENIZATION OF A MODEL OF COMPRESSIBLE MISCIBLE FLOW IN POROUS MEDIA**

## **HOMOGÉNÉISATION D' UN MODÈLE D' ÉCOULEMENTS MISCIBLES COMPRESSIBLES EN MILIEU POREUX**

**Y. AMIRAT <sup>1</sup>**

**K. HAMDACHE <sup>2</sup>**

**A. ZIANI <sup>3</sup>**

---

<sup>1</sup> *INRIA, Domaine de Voluceau, Rocquencourt, B.P. 105 F-78135 LE CHESNAY Cedex*

<sup>2</sup> *ENSTA-GHN, Centre de l'Yvette, Chemin de la Hunière, F-91120 PALAISEAU*

<sup>3</sup> *Institut de Mathématiques, USTHB, B.P. 32 EL ALIA DZ-16111, ALGER*

## **Abstract**

We consider a 1-D model of flow of two miscible fluids with same compressibility factor in an heterogeneous porous media. We study the existence and homogenization of the parabolic-hyperbolic system governing this flow and show the stability of the derivation of the macroscopic level .

## **Résumé**

On considère un modèle 1-D d'écoulements de deux fluides miscibles ayant même facteur de compressibilité. On étudie l'existence et l'homogénéisation du système parabolique-hyperbolique gouvernant de tels écoulements, et on montre que le problème homogénéisé est de même type.

## **Key-words**

Characteristics, Homogenization, Hyperbolic, Oscillations, Parabolic, Porous Media.

## **Mots-Clés**

Caractéristiques, Homogénéisation, Hyperbolique, Milieu Poreux, Oscillations, Parabolique.

## 0. Introduction

With the work of DARCY [10] starts the theory of fluid flow in porous media. From the description he gave of the relationship between pressure and flux of a fluid in a homogeneous porous media, several attempts to generalize the concept to multiphase flow have been made, we refer to BEAR [5], HOUPEURT [13], MATHERON [19], MARLES [18], MUSKAT [21], SCHEIDEGGER [24] and SLATTERY [26].

The objective to give a clear formulation and mathematical description is not achieved considering the complexity of phenomena occurring in a media in presence of a mixture of several fluids and solids.

To derive a macroscopic model for different flows in a heterogeneous porous media various tools have been used, such as the averaging methods and theory of homogenization which are developed in AURIAULT-SANCHEZ PALENCIA [3], BAKHVALOV-PANASENKO [4], MARCHENKO-HROUSLOV [16] and SANCHEZ PALENCIA [23]. Another approach using stochastic analysis and averaging using Markov processes lead to some results in MATHERON [19], CAFLISCH-RUBINSTEIN [8], RUBINSTEIN [22], FIGARI-PAPANICOLAOU-RUBINSTEIN [12] and SHVIDLER [25].

This paper deals with a simpler model in one dimension of the flow of two miscible fluids in a heterogeneous porous media. The aim is to derive a set of effective equation from the microscopic level.

One 1-D non linear model for incompressible miscible displacement has been considered in [1], we have proved the stability of the Darcy law with respect to the oscillations of some parameters.

The homogenization of the first order, linear hyperbolic equation studied in [2] induces for the limiting equation satisfied by the concentration of the fluid some memory effects due to dispersion phenomena. In what follows, we consider two miscible fluids. The composition of the mixture is given by the mass concentration. We neglect the molecular diffusion and dispersion and omit the gravitational terms. The viscosity is assumed constant and the fluids have the same constant compressibility factor. The porosity  $\phi$  and permeability  $k$  of the rock are oscillating functions of the spatial variable  $x$  and depend on a parameter  $\epsilon > 0$  connected with the microstructure.

If the functions  $p, q$  and  $u$  stand for pressure, rate of flow and concentration of mass respectively, the flow is governed by the following system of parabolic-hyperbolic type :

( see CHAVENT-JAFFRE [9], DOUGLAS-ROBERTS [11])

$\Omega = ]0, 1[, T$  is a fixed positive number and  $\Omega_T = ]0, 1[ \times \Omega$

$(p^\epsilon, q^\epsilon, u^\epsilon)$  satisfies :

$$(P_\epsilon) \begin{cases} \phi^\epsilon(x) \frac{\partial}{\partial t} p^\epsilon(t, x) + \frac{\partial}{\partial x} q^\epsilon(t, x) = 0 \\ q^\epsilon(t, x) = -k^\epsilon(x) \frac{\partial}{\partial x} p^\epsilon(t, x) & (t, x) \in \Omega_T \\ q^\epsilon(t, 0) = q_1^\epsilon(t), \quad q^\epsilon(t, 1) = q_2^\epsilon(t) & t \in ]0, T[ \\ p^\epsilon(0, x) = p_0^\epsilon(x) & x \in \Omega \end{cases}$$

$$(H_\epsilon) \begin{cases} \phi^\epsilon(x) \frac{\partial}{\partial t} u^\epsilon(t, x) + q^\epsilon(t, x) \frac{\partial}{\partial x} u^\epsilon(t, x) = 0 & (t, x) \in \Omega_T \\ 0 \leq u^\epsilon(t, x) \leq 1 \\ u^\epsilon(t, 0) = u_1^\epsilon(t), \quad 0 \leq u_1^\epsilon(t) \leq 1 & t \in ]0, T[ \\ u^\epsilon(0, x) = u_0^\epsilon(x), \quad 0 \leq u_0^\epsilon(x) \leq 1 & x \in \Omega \end{cases}$$

The boundary data  $q_1^\epsilon, q_2^\epsilon, u_1^\epsilon$  and the initial data  $p_0^\epsilon$  and  $u_0^\epsilon$  are taken dependent on  $\epsilon$ , this dependence will be specified further. The boundary condition for  $u^\epsilon$  is given for  $x = 0$  because of the positivity of  $q^\epsilon$  as will be seen.

We want to analyse the influence of heterogeneities - characterized by the parameter  $\epsilon$  - on the effective behavior of the pressure, flux and mass concentration.

The summary of the paper is the following :

In section 1 we consider the case of a periodic microstructure and derive formally the equations obtained asymptotically for the pressure and concentration. In section 2 we recall some properties concerning a parabolic equation with discontinuous coefficients and study the behavior of the flux  $q^\epsilon$  and derive the homogenized equation for  $q$ . The third section is devoted to the existence of a weak solution for the problem  $(H^\epsilon)$  and in section 4 we look for the limiting equation for  $u^\epsilon$ .

In a forthcoming paper we will discuss the more realistic model where the two fluids have different compressibility factors.

## 1. Asymptotic study of media with periodic microstructure

In this section we use the asymptotic methods introduced by BENSOUSSAN- LIONS- PAPANICOLAOU [6], LIONS [15] and SANCHEZ PALENCIA [23] to derive the limiting equations in the case of a periodic structure, namely we assume that the coefficients  $\phi^\epsilon$  and  $k^\epsilon$  are given by :

$$(1.1) \quad \begin{cases} \phi^\epsilon(x) = \phi(x, \frac{x}{\epsilon}), \\ k^\epsilon(x) = k(x, \frac{x}{\epsilon}), \end{cases}$$

where the functions  $\phi(x, y)$  and  $k(x, y)$  are 1-periodic with respect to the  $y$  variable .

We look for the pressure  $p^\epsilon(t, x)$  in the form of an asymptotic expansion :

$$p^\epsilon(t, x) = p^0(t, x, y) + \epsilon p^1(t, x, y) + \epsilon^2 p^2(t, x, y) + \dots$$

where  $y = \frac{x}{\epsilon}$ , and  $p^i(t, x, y)$  is a periodic functions of the variables  $y$  for  $i = 0, 1, \dots$

Using the equation :

$$(1.2) \quad \phi^\epsilon(x) \frac{\partial}{\partial t} p^\epsilon(t, x) - \frac{\partial}{\partial x} (k^\epsilon(x) \frac{\partial}{\partial x} p^\epsilon(t, x)) = 0$$

and expanding we obtain the following relations :

$$(1.3) \quad \begin{cases} \frac{\partial}{\partial y}(k(x, y) \frac{\partial}{\partial y} p^0) = 0, \\ \frac{\partial}{\partial y}(k(x, y) \frac{\partial}{\partial y} p^1) + \frac{\partial}{\partial y}(k(x, y) \frac{\partial}{\partial x} p^0) + \frac{\partial}{\partial x}(k(x, y) \frac{\partial}{\partial y} p^0) = 0, \\ \phi(x, y) \frac{\partial}{\partial t} p^0 - \frac{\partial}{\partial y}(k(x, y) \frac{\partial}{\partial y} p^2) \\ = \{ \frac{\partial}{\partial y}(k(x, y) \frac{\partial}{\partial x} p^0) + \frac{\partial}{\partial y}(k(x, y) \frac{\partial}{\partial x} p^1) + \frac{\partial}{\partial x}(k(x, y) \frac{\partial}{\partial y} p^1) \}, \\ \text{etc} \dots \end{cases}$$

The first one implies that  $p^0$  doesn't depend on  $y$  so  $p^0 = p^0(t, x)$ . The second one implies that  $p^1$  is well known :

$$p^1(t, x, y) = \chi(x, y) \frac{\partial}{\partial x} p^0(t, x) + \tilde{p}^1(t, x),$$

where  $\chi$  is the unique solution of :

$$(1.4) \quad \begin{cases} -\frac{\partial}{\partial y}(k(x, y) \frac{\partial}{\partial y} \chi) = \frac{\partial}{\partial y} k(x, y), \\ \chi \text{ 1- periodic on } y \text{ and } \int_0^1 \chi(x, y) dy = 0. \end{cases}$$

The periodicity of  $p^2$  leads to the following homogenized equation :

$$(1.5) \quad \tilde{\phi}(x) \frac{\partial}{\partial t} p^0 - \frac{\partial}{\partial x} \left\{ \int_0^1 k(x, y) \left( 1 + \frac{\partial}{\partial y} \chi(x, y) \right) dy \frac{\partial}{\partial x} p^0 \right\} = 0.$$

where  $\tilde{\phi}(x) = \int_0^1 \phi(x, y) dy$ . If we assume that the data are independent of  $\epsilon$  then  $p^0$  is well defined and also  $p^l(t, x, y)$   $l \geq 1$ .

Therefore the flux  $q^\epsilon$  is given by :

$$(1.6) \quad q^\epsilon(t, x) = q^0(t, x, y) + \epsilon q^1(t, x, y) + \epsilon^2 q^2(t, x, y) + \dots$$

where  $q^l(t, x, y) = -k(x, y) \left[ \frac{\partial}{\partial x} p^l + \frac{\partial}{\partial y} p^{l+1} \right]$   $l \geq 0$ .

As for the pressure we assume that the concentration  $u^\epsilon$  also admits the following asymptotic expansion in  $\epsilon$  :

$$(1.7) \quad u^\epsilon(t, x) = u^0(t, x, y) + \epsilon u^1(t, x, y) + \epsilon^2 u^2(t, x, y) + \dots$$

Plugging (1.6) and (1.7) into the transport equation :

$$(1.8) \quad \phi^\epsilon(x) \frac{\partial}{\partial t} u^\epsilon + q^\epsilon(t, x) \frac{\partial}{\partial x} u^\epsilon = 0.$$

we get the following system of equations

$$(1.9) \quad \begin{cases} q^0(t, x, y) \frac{\partial}{\partial y} u^0 = 0, \\ \phi(x, y) \frac{\partial}{\partial t} u^0 + q^0(t, x, y) \left[ \frac{\partial}{\partial x} u^0 + \frac{\partial}{\partial y} u^1 \right] = 0 \\ \text{etc} \dots \end{cases}$$

As  $q^0 \neq 0$  the first equation implies  $u^0 = u^0(t, x)$ , the second one can be written as

$$(1.10) \quad \frac{\phi(x, y)}{q^0(t, x, y)} \frac{\partial}{\partial t} u^0 + \frac{\partial}{\partial x} u^0 = - \frac{\partial}{\partial y} u^1.$$

Using the periodicity of  $u^1$  in  $y$  and integrating over the interval  $(0, 1)$  we find :

$$(1.11) \quad \int_0^1 \frac{\phi(x, y)}{q^0(t, x, y)} dy \frac{\partial}{\partial t} u^0 + \frac{\partial}{\partial x} u^0 = 0$$

Taking into account the value of  $q^0$  in terms of  $p^0$

$$q^0(t, x, y) = -k(x, y) \left[ 1 + \frac{\partial}{\partial y} \chi \right] \frac{\partial}{\partial x} p^0$$

we get the macroscopic equation for the concentration :

$$(1.12) \quad \int_0^1 \frac{\phi(x, y)}{k(x, y) \left[ 1 + \frac{\partial}{\partial y} \chi \right]} dy \frac{\partial}{\partial t} u^0 + \frac{\partial}{\partial x} p^0 \frac{\partial}{\partial x} u^0 = 0.$$

So we have proved, in this setting that the homogenized equations (1.5) and (1.12) have the same form as the microscopic one (1.2), (1.8).

**Remark 1.1**

If the data  $p^\epsilon(x, 0) = p_0^\epsilon(x)$  satisfies :

$$(1.13) \quad -\frac{\partial}{\partial x} \left( k^\epsilon \frac{\partial}{\partial x} p_0^\epsilon \right) = g \quad \text{with} \quad g \in L^2(\Omega),$$

then we will see later in §2 that the pressure  $p^\epsilon$  is such that

$$\frac{\partial}{\partial t} p^\epsilon \in L^2(\Omega_T),$$

and therefore we obtain :

$$q^\epsilon(t, x) \rightarrow \tilde{q}^0(t, x) = - \int_0^1 k(x, y) \left[ 1 + \frac{\partial}{\partial y} \chi \right] dy \frac{\partial}{\partial x} p^0 \quad \text{in } L^2(\Omega_T).$$

(1.6) is then replaced by



$$(1.6') \quad q^\epsilon(t, x) = \tilde{q}^0(t, x) + \epsilon q^1(t, x, y) + \epsilon^2 q^2(t, x, y) + \dots$$

hence (1.12) by

$$(1.12') \quad \tilde{\phi}(x) \frac{\partial}{\partial t} u^0 + \tilde{q}^0(t, x) \frac{\partial}{\partial x} u^0 = 0.$$

The case where the initial data doesn't fulfill such a matching condition leads to a boundary layer (see [23]).

## 2. Homogenization of a problem of parabolic type

In the sequel we consider the problem :

$$(2.1) \quad \begin{cases} \phi^\epsilon(x) \frac{\partial}{\partial t} p^\epsilon(t, x) + \frac{\partial}{\partial x} q^\epsilon(t, x) = 0, \\ q^\epsilon(t, x) = -k^\epsilon(x) \frac{\partial}{\partial x} p^\epsilon(t, x) \quad (t, x) \in \Omega_T, \\ q^\epsilon(0, t) = q_1^\epsilon(t), \quad q^\epsilon(1, t) = q_2^\epsilon(t) \quad t \in ]0, T[, \\ p^\epsilon(x, 0) = p_0^\epsilon(x) \quad x \in \Omega, \end{cases}$$

with the assumptions :

$$(2.2) \quad \begin{cases} \phi^\epsilon, k^\epsilon \in L^\infty(\Omega) \\ 0 < \phi_- \leq \phi^\epsilon \leq \phi^+, \quad 0 < k_- \leq k^\epsilon \leq k^+. \end{cases}$$

$$(2.3) \quad \begin{cases} \text{There exist a constant } C > 0 \text{ such that for all } \epsilon > 0 \\ \|q_i^\epsilon\|_{W^{1,\infty}(0,T)} \leq C \quad i = 1, 2 \quad \|p_0^\epsilon\|_{H^1(\Omega)} \leq C \end{cases}$$

The homogenization of probleme of type (2.1) has been treated by SPAGNOLO [27] in the framework of **G**-convergence and also by MARKOV-OLEINIK [17] in a more general setting with the same motivation to investigate the properties of disperse media.

As the flux  $q^\epsilon$  appears as a coefficient in the transport equation it is important to point out more precisely its properties for the hypothesis (2.2),(2.3).

Since

$$\frac{\partial}{\partial x} p^\epsilon = -\frac{1}{k^\epsilon} q^\epsilon \quad \text{and} \quad \frac{\partial}{\partial t} p^\epsilon = -\frac{1}{\phi^\epsilon} \frac{\partial}{\partial x} q^\epsilon$$

we can formulate a problem satisfied by  $q^\epsilon$  :

$$(2.4) \quad \begin{cases} \frac{1}{k^\epsilon} \frac{\partial}{\partial t} q^\epsilon - \frac{\partial}{\partial x} \left[ \frac{1}{\phi^\epsilon(x)} \frac{\partial}{\partial x} q^\epsilon \right] = 0 \quad \text{in } \Omega_T, \\ q^\epsilon(t, 0) = q_1^\epsilon(t), \quad q^\epsilon(t, 1) = q_2^\epsilon(t), \quad t \in ]0, T[, \\ q^\epsilon(0, x) = -k^\epsilon(x) \frac{\partial}{\partial x} p_0^\epsilon(x) \equiv q_0^\epsilon(x) \quad x \in \Omega. \end{cases}$$

We begin to render the boundary condition homogeneous by introducing for  $T > 0$  the elliptic problem :

$$(2.5) \quad \begin{cases} -\frac{\partial}{\partial x} \left( \frac{1}{\phi^\epsilon} \frac{\partial}{\partial x} Q^\epsilon \right) = 0 & \text{in } \Omega \\ Q^\epsilon(0, t) = q_1^\epsilon(t), \quad Q^\epsilon(1, t) = q_2^\epsilon(t), & t \in ]0, T[. \end{cases}$$

The solution is given by :

$$(2.6) \quad q^\epsilon(t, x) = q_1^\epsilon(t) + \frac{\Psi^\epsilon(x)}{\Psi^\epsilon(1)} [q_2^\epsilon(t) - q_1^\epsilon(t)]$$

where  $\Psi^\epsilon(x) = \int_0^x \phi^\epsilon(\xi) d\xi$  Because of (2.2) and (2.3), we have :

$$(2.7) \quad Q^\epsilon \text{ belongs to a bounded subset of } W^{1,\infty}(\Omega_T).$$

The function  $v^\epsilon$  defined by :

$$(2.8) \quad v^\epsilon = q^\epsilon - Q^\epsilon$$

is such that :

$$(2.9) \quad \begin{cases} \frac{1}{k^\epsilon(x)} \frac{\partial}{\partial t} v^\epsilon - \frac{\partial}{\partial x} \left[ \frac{1}{\phi^\epsilon(x)} \frac{\partial}{\partial x} v^\epsilon \right] = f^\epsilon \equiv -\frac{1}{k^\epsilon} \frac{\partial}{\partial t} Q^\epsilon \\ v^\epsilon(0, t) = v^\epsilon(1, t) = 0 \quad t \in ]0, T[ \\ v^\epsilon(x, 0) = v_0^\epsilon(x) \equiv q_0^\epsilon(x) - Q^\epsilon(x, 0) \quad x \in \Omega \end{cases}$$

$v^\epsilon$  is a solution of a non-degenerate equation of parabolic type with source term  $f^\epsilon$  uniformly bounded in  $L^\infty(\Omega_T)$  and initial data bounded in  $L^2(\Omega)$ .

The standard theory of linear parabolic equations (cf LADYZENSKAJA, SOLONNIKOV, URAL'CEVA [14]) shows that for all  $T > 0$ ,  $v^\epsilon$  exists and is unique in the space  $L^\infty(0, T; L^2(\Omega)) \cap L^2(0, T; H_0^1(\Omega))$ .

The classical energy estimate

$$\frac{1}{2} \frac{d}{dt} \int_\Omega \frac{1}{k^\epsilon(x)} |v^\epsilon(t, x)|^2 dx + \int_\Omega \frac{1}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial x} v^\epsilon(t, x) \right|^2 dx = \int_\Omega f^\epsilon(t, x) v^\epsilon(t, x) dx$$

shows that thanks to (2.2), (2.3) and (2.7), the sequence  $\{v^\epsilon\}_{\epsilon>0}$  is uniformly bounded with respect to  $\epsilon$  in  $L^\infty(0, T; L^2(\Omega)) \cap L^2(0, T; H_0^1(\Omega))$ .

The solution  $v^\epsilon$  enjoys the following property

### Lemma 2.1

Under the hypothesis (2.2), (2.3) the solution  $v^\epsilon$  of (2.9) has the regularity :

$$(2.10) \quad t \frac{\partial}{\partial t} v^\epsilon \text{ belongs to a bounded subset of } L^2(\Omega_T)$$

To this end we multiply the equation (2.9) by  $t^2 \frac{\partial}{\partial t} v^\epsilon$  and integrate over  $\Omega$  therefore we have

$$\int_{\Omega} \frac{t^2}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial t} v^\epsilon(t, x) \right|^2 dx + \frac{1}{2} \int_{\Omega} t^2 \frac{1}{\phi^\epsilon(x)} \frac{\partial}{\partial t} \left| \frac{\partial}{\partial x} v^\epsilon(t, x) \right|^2 dx = \int_{\Omega} t^2 f^\epsilon(t, x) \frac{\partial}{\partial t} v^\epsilon(t, x) dx$$

so

$$t^2 \frac{1}{\phi^\epsilon(x)} \frac{\partial}{\partial t} \left| \frac{\partial}{\partial x} v^\epsilon(t, x) \right|^2 = \frac{\partial}{\partial t} \left\{ t^2 \frac{1}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial x} v^\epsilon(t, x) \right|^2 \right\} - 2t \frac{1}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial x} v^\epsilon(t, x) \right|^2$$

Integration in time on  $(0, T)$  yields :

$$\begin{aligned} & \int_{\Omega_T} \frac{t^2}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial t} v^\epsilon(t, x) \right|^2 dx dt + \frac{T^2}{2} \int_{\Omega} \frac{1}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial x} v^\epsilon(t, x) \right|^2 dx \\ &= \int_{\Omega_T} t^2 f^\epsilon(t, x) \frac{\partial}{\partial t} v^\epsilon(t, x) dx dt + \int_{\Omega_T} \frac{t}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial x} v^\epsilon(t, x) \right|^2 dx dt, \end{aligned}$$

and  $\forall \eta > 0, \exists C_\eta > 0$  such that the left hand side is less than

$$T \int_{\Omega_T} \frac{1}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial t} v^\epsilon(t, x) \right|^2 dx dt + \eta \int_{\Omega_T} t^2 \left| \frac{\partial}{\partial t} v^\epsilon(t, x) \right|^2 dx dt + C_\eta \int_{\Omega_T} t^2 |f^\epsilon(t, x)|^2 dx dt$$

which implies with (2.2) and (2.3) that the sequence  $\{v^\epsilon\}_{\epsilon>0}$  satisfies :

$$\left\| t \frac{\partial}{\partial t} v^\epsilon \right\|_{L^2(\Omega_T)} \leq C$$

where  $C$  is a constant independent of  $\epsilon$ .

### Remark 2.2

From (2.9) and (2.10) we deduce that for all  $\delta > 0$  :

$$\frac{\partial}{\partial x} \left[ \frac{1}{\phi^\epsilon} \frac{\partial}{\partial x} v^\epsilon \right] \in L^2([\delta, T] \times \Omega),$$

and then

$$\left[ \frac{1}{\phi^\epsilon} \frac{\partial}{\partial x} v^\epsilon \right] \in L^2([\delta, T[, H_0^1(\Omega)),$$

so  $\frac{\partial}{\partial x} v^\epsilon$  belongs to a bounded set of  $L^2(\delta, T; L^\infty(\Omega))$ , which with (2.7) implies that :

$$(2.11) \quad \left\| \frac{\partial}{\partial x} q^\epsilon \right\|_{L^2(\delta, T; L^\infty(\Omega))} \leq C,$$

where  $C$  is a constant independent of  $\epsilon$ .

To obtain an estimate of  $\frac{\partial}{\partial t}v^\epsilon$  in  $L^2(\Omega_T)$ , we actually need an additional condition on the data  $p_0^\epsilon$ . Indeed, from (2.9) we get, after multiplication by  $\frac{\partial}{\partial t}v^\epsilon$  and integration over  $\Omega$  and from 0 to  $t$ ,

$$\int_{\Omega} \frac{1}{k^\epsilon(x)} \left| \frac{\partial}{\partial t}v^\epsilon(t, x) \right|^2 dx + \frac{1}{2} \frac{d}{dt} \int_{\Omega} \frac{1}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial x}v^\epsilon(t, x) \right|^2 dx = \int_{\Omega} f^\epsilon(t, x) \frac{\partial}{\partial t}v^\epsilon(t, x) dx$$

$$\begin{aligned} & \int_0^t \int_{\Omega} \frac{1}{k^\epsilon(x)} \left| \frac{\partial}{\partial t}v^\epsilon(\sigma, x) \right|^2 dx d\sigma + \frac{1}{2} \int_{\Omega} \frac{1}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial x}v^\epsilon(t, x) \right|^2 dx \\ &= \int_{\Omega} f^\epsilon(\sigma, x) \frac{\partial}{\partial t}v^\epsilon(\sigma, x) dx d\sigma + \frac{1}{2} \int_{\Omega} \frac{1}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial x}v^\epsilon(0, x) \right|^2 dx \end{aligned}$$

Making use of (2.4) (2.9) (2.2) and (2.3) we have :

$$\frac{\partial}{\partial x}v^\epsilon(0, x) = -\frac{\partial}{\partial x}(k^\epsilon(x) \frac{\partial}{\partial x}p_0^\epsilon) - \frac{\partial}{\partial x}Q^\epsilon(0, x).$$

In order that the term

$$\int_{\Omega} \frac{1}{\phi^\epsilon(x)} \left| \frac{\partial}{\partial x}v^\epsilon(0, x) \right|^2 dx$$

be uniformly bounded with respect to  $\epsilon$  it is sufficient that  $p_0^\epsilon$  satisfies an elliptic problem

$$(2.12) \quad -\frac{\partial}{\partial x}(k^\epsilon(x) \frac{\partial}{\partial x}p_0^\epsilon) = g^\epsilon$$

for any sequence with  $\|g^\epsilon\|_{L^2(\Omega_T)} \leq C$ . Consequently we obtain :

$$(2.13) \quad \left\| \frac{\partial}{\partial t}v^\epsilon \right\|_{L^2(\Omega_T)} \leq C,$$

where  $C$  is a constant independent of  $\epsilon$ .

Let

$$\gamma^\epsilon = \min \left\{ \min_{t \in ]0, T[} q_1^\epsilon(t), \min_{t \in ]0, T[} q_2^\epsilon(t), \min_{x \in \Omega} q_0^\epsilon(x) \right\}$$

$$\Gamma^\epsilon = \max \left\{ \max_{t \in ]0, T[} q_1^\epsilon(t), \max_{t \in ]0, T[} q_2^\epsilon(t), \max_{x \in \Omega} q_0^\epsilon(x) \right\}$$

Using the maximum principle (cf [14] ) we have :

**Lemma 2.3**

The solution  $q^\epsilon$  of (2.4) verifies the following property :

$$(2.14) \quad \min\{\gamma^\epsilon, 0\} \leq q^\epsilon(t, x) \leq \max\{0, \Gamma^\epsilon\}$$

for almost all  $(t, x) \in \Omega_T$

If  $\gamma^\epsilon \geq \gamma > 0$  then  $q^\epsilon(t, x) > 0$  which implies that the hyperbolic problem  $(H_\epsilon)$  is well posed for the boundary condition at  $x = 0$ .

**Remark 2.4**

The hypotheses (2.2) (2.3) imply that we have the energy estimate for pressure  $p^\epsilon$  :

$$(2.15) \quad \left\| \frac{\partial}{\partial t} p^\epsilon \right\|_{L^2(\Omega_T)} \leq C$$

where  $C$  is a constant independent of  $\epsilon$ .

we have also :

$$(2.16) \quad \left\| \frac{\partial^2}{\partial t^2} p^\epsilon \right\|_{L^2(\Omega_T^\epsilon)} \leq C, \quad \left\| \frac{\partial^2}{\partial t \partial x} p^\epsilon \right\|_{L^2(\Omega_T^\epsilon)} \leq C,$$

where  $C$  is a constant independent of  $\epsilon$  and  $\Omega_T^\delta = ]\delta, T[ \times \Omega$ .

Assume that

$$(2.17) \quad \phi^\epsilon \rightarrow \phi, \quad \frac{1}{k^\epsilon} \rightarrow \frac{1}{k} \quad \text{in } L^\infty(\Omega) \text{ weak } *.$$

From (2.3) we deduce :

$$q_i^\epsilon \rightarrow q_i \quad \text{in } L^\infty(O, T) \quad i = 1, 2,$$

$$q_0^\epsilon \rightharpoonup q_0 \quad \text{in } L^2(\Omega) \quad \text{weakly}.$$

Using the techniques introduced by MURAT [20], TARTAR [28] and the result of MARKOV-OLEINIK [17], we obtain the :

**Theorem 2.5**

The sequence  $\{v^\epsilon\}_{\epsilon>0}$  of solutions of (2.4) is such that :

$$(2.18) \quad \begin{cases} q^\epsilon \rightharpoonup q & \text{in } L^2(0, T; L^2(\Omega)) \text{ weakly,} \\ q^\epsilon \rightarrow q & \text{uniformly in } [\delta, T] \times \Omega \quad \forall \delta > 0 \end{cases}$$

where  $q$  is the unique solution of :

$$(2.19) \quad \begin{cases} \frac{1}{k(x)} \frac{\partial}{\partial t} q - \frac{\partial}{\partial x} \left( \frac{1}{\phi(x)} \frac{\partial}{\partial x} q \right) = 0 & \text{in } \Omega_T, \\ q(t, 0) = q_1(t), \quad q(t, 1) = q_2(t) & t \in (0, T), \\ q(0, x) = q_0(x) & x \in \Omega. \end{cases}$$

### Corollary 2.6

The solution  $q$  of (2.19) belongs to  $C([0, T]; L^2(\Omega)) \cap L^2(0, T; H^1(\Omega))$  and satisfies

$$\begin{aligned} \|t \frac{\partial}{\partial t} v^\epsilon\|_{L^2(\Omega_T)} &\leq C, \\ \gamma &\leq q(t, x) \leq \Gamma \quad \text{a.e. } (t, x) \in \Omega_T, \\ \text{where } 0 &< \gamma \leq \gamma^\epsilon \leq \Gamma^\epsilon \leq \Gamma. \end{aligned}$$

### Remark 2.7

If the sequence  $\{p_0^\epsilon\}_{\epsilon>0}$  satisfies (2.12) and

$$g^\epsilon \rightharpoonup g \quad \text{in } L^2(\Omega) \quad \text{weakly}$$

then :

$$q_0^\epsilon \rightharpoonup q_0 = -k \frac{\partial}{\partial x} p_0 \quad \text{in } L^2(\Omega) \quad \text{weakly}$$

where  $k$  is defined by (2.17) and  $p_0$  is the solution of the homogenized elliptic problem (2.13) described by :

$$-\frac{\partial}{\partial x} \left( k \frac{\partial}{\partial x} p_0 \right) = g \quad \text{in } L^2(\Omega).$$

## 3. Study of a hyperbolic first order equation in a non-divergence form

The function  $q^\epsilon$  being defined by (2.4), we want to investigate weak solutions for the problem :

$$(3.1) \quad \begin{cases} \phi^\epsilon(x) \frac{\partial}{\partial t} u^\epsilon(t, x) + q^\epsilon(t, x) \frac{\partial}{\partial x} u^\epsilon(t, x) = 0 & \text{in } \Omega_T, \\ u^\epsilon(t, 0) = u_1^\epsilon(t), \quad t \in ]0, T[, \\ u^\epsilon(0, x) = u_0^\epsilon(x) \quad x \in \Omega, \\ 0 \leq u^\epsilon(t, x) \leq 1 \quad (t, x) \in \Omega_T. \end{cases}$$

As in [1], we introduce the following notion of a weak solution :

**Definition 3.1**

A function  $u^\epsilon$  belonging to  $L^\infty(\Omega_T)$  is a weak solution of (3.1) if  $u^\epsilon$  has its values in  $[0,1]$  and satisfies :

$$(3.2) \quad \begin{aligned} & \int_0^T \int_\Omega \left\{ \phi^\epsilon \frac{\partial}{\partial t} \varphi + q^\epsilon(t, x) \frac{\partial}{\partial x} \varphi + \frac{\partial}{\partial x} q^\epsilon(t, x) \varphi \right\} u^\epsilon(t, x) dx dt \\ &= - \int_\Omega \phi^\epsilon(x) u_0^\epsilon(x) \varphi(0, x) dx - \int_0^T q_1^\epsilon(t) u_1^\epsilon(t) \varphi(t, 0) dt \end{aligned}$$

for any function  $\varphi \in C_0^1([0, T] \times [0, 1])$ .

To define such a solution we begin by the study of characteristic curves  $X^\epsilon$  associated with the first order differential operator :

$$(3.3) \quad \phi^\epsilon(x) \frac{\partial}{\partial t} + q^\epsilon(t, x) \frac{\partial}{\partial x}.$$

For  $\epsilon > 0$ , we consider the equation of the characteristic curve associated to (3.3)

$$(3.4) \quad \begin{cases} \frac{\partial}{\partial t} X^\epsilon = W^\epsilon(t, x) \equiv \frac{q^\epsilon(t, X^\epsilon)}{\phi^\epsilon(X^\epsilon)} \\ X^\epsilon(s) = x, \quad s \in [0, T], \quad x \in \Omega. \end{cases}$$

The function  $W^\epsilon(t, x)$  doesn't satisfy the Caratheodory conditions for the existence of a solution  $X^\epsilon$ , we proceed by regularization of coefficients  $\phi^\epsilon$  and  $q^\epsilon$  of (3.3).

For the remainder of this section we drop the subscript  $\epsilon$ .

Let  $\phi^n$  and  $q^n$  two sequences of regular functions such that :

$$(3.5) \quad \begin{cases} \phi^n \rightarrow \phi \quad \text{in } L^2(\Omega), \\ q^n \rightarrow q \quad \text{in } L^2(\Omega_T), \\ \phi^n \in C^1(\Omega) \quad \phi_- \leq \phi^n(x) \leq \phi^+, \\ 0 < \gamma \leq q^n(t, x) \leq \Gamma \quad (t, x) \in \Omega_T, \\ \left\| \frac{\partial}{\partial x} q^n(t, \cdot) \right\|_{L^\infty(\Omega)} \leq a(t) \quad a \in L^2(0, T) \quad \text{independent on } n. \end{cases}$$

For each integer  $n$  we introduce  $X^n$  uniquely defined by

$$(3.6) \quad \begin{cases} \frac{\partial}{\partial t} X^n = \frac{q^n(t, X^n)}{\phi^n(X^n)}, \\ X^n(s) = x, \quad s \in [0, T], \quad x \in \Omega. \end{cases}$$

If  $\Psi^n$  is the antiderivative of  $\phi^n$  given by

$$(3.7) \quad \Psi^n(x) = \int_0^x \phi^n(\xi) d\xi,$$

the equation (3.6) can also be written

$$(3.8) \quad \frac{\partial}{\partial t} \Psi^n(X^n) = q^n(t, X^n),$$

which gives after integration the following *weak formulation* of the characteristic curve (3.6)

$$(3.9) \quad \Psi^n(X^n) = \int_s^t q^n(\sigma, X^n(\sigma; x, s)) d\sigma + \Psi^n(x).$$

Now, we want to derive some properties of the sequence  $\{X^n\}_{n \geq 0}$ .

First the estimate :

$$(3.10) \quad \left\| \frac{\partial}{\partial t} X^n \right\|_{L^\infty(\Omega_T)} \quad \text{is valid.}$$

Differentiate (3.9) with respect to  $x$ , to obtain the following relation :

$$(3.11) \quad \phi^n(X^n) \frac{\partial}{\partial x} X^n = \int_s^t \frac{\partial}{\partial x} q^n(\sigma, X^n(\sigma; x, s)) d\sigma + \phi^n(x).$$

We set

$$w^n(t) = \sup_{x \in \Omega} \left| \frac{\partial}{\partial x} X^n(t; x, s) \right|.$$

From the hypothesis (3.5) we deduce the inequality :

$$(3.12) \quad \phi_- w^n(t) \leq \int_s^t a(\sigma) w^n(\sigma) d\sigma + \phi^+,$$

whence by the Gronwall lemma :

$$(3.13) \quad w^n(t) \leq b(t) \quad \text{with} \quad b \in L^2(0, T),$$

i.e.

$$(3.14) \quad \left\| \frac{\partial}{\partial x} X^n \right\|_{L^2(0, T; L^\infty(\Omega))} \leq C.$$

Also we define implicitly  $S^n(t, x)$  by :



$$(3.15) \quad X^n(S^n(t, x); x, t) = 0, (t, x) \in \Omega_T,$$

which implies

$$(3.16) \quad \phi^n(X^n) \frac{\partial}{\partial x} X^n = \int_t^{S^n(t, x)} q^n(\sigma, X^n(\sigma; x, s)) d\sigma + \Psi^n(x)$$

From (3.15), we have using (3.14) that

$$(3.17) \quad \left\| \frac{\partial}{\partial x} S^n \right\|_{L^2(0, T; L^\infty(\Omega))} \leq C, \quad \left\| \frac{\partial}{\partial t} X^n \right\|_{L^\infty(\Omega_T)} \leq C.$$

Hence, there exist two subsequences, also denoted by  $X^n, S^n$  such that

$$(3.18) \quad X^n \rightarrow X, \quad S^n \rightarrow S \quad \text{strongly in } L^2(\Omega_T).$$

Given the regularity properties (3.5) of the coefficients, the function  $u^n$  defined by

$$(3.19) \quad u^n(t, x) = \begin{cases} u_0^\epsilon(X^n(0; x, t)), & \text{if } x > X^n(t; 0, 0) \\ u_1^\epsilon(S^n(t, x)), & \text{otherwise,} \end{cases}$$

is the classical solution of the problem

$$(3.20) \quad \begin{cases} \phi^n \frac{\partial}{\partial t} u^n(t, x) + q^n(t, x) \frac{\partial}{\partial x} u^n(t, x) = 0 & \text{in } \Omega_T, \\ u^n(t, 0) = u_1^\epsilon(t) & t \in ]0, T[, \\ u^n(0, x) = u_0^\epsilon(x) & x \in \Omega. \end{cases}$$

Clearly

$$0 \leq u^n(t, x) \leq 1, \quad (t, x) \in \Omega_T$$

if the data satisfy

$$0 \leq u_0^\epsilon, u_1^\epsilon \leq 1.$$

For the limiting functions  $X$  and  $S$ , we establish some necessary a-priori estimates independent of  $\epsilon$  in order to pass to the limit as  $\epsilon \rightarrow 0$  in the next section

**Lemma 3.2**

$\forall \epsilon > 0$ ,  $X$  and  $S$  satisfy the equations

$$(3.21) \quad \begin{cases} \Psi(X) = \int_s^t q(\sigma, X(\sigma; x, s)) d\sigma + \Psi(x), \\ X(S(t, x); x, t) = 0 \quad \text{a.e. in } \Omega_T. \end{cases}$$

**Lemma 3.3**

$X$  and  $S$  belong to  $V_\delta$  for all  $\delta > 0$  where

$$(3.22) \quad V_\delta = \{v \in L^\infty(\Omega_T), \frac{\partial}{\partial t}v \in L^\infty(\Omega_T), \frac{\partial}{\partial x}v \in L^2(\delta, T; L^\infty(\Omega))\}.$$

**Proof.** It is based on the above procedure and on the property (2.11) of  $q^\epsilon$  and hypothesis (2.2).

Let us define the function  $u$  by

$$(3.23) \quad u(t, x) = \begin{cases} u_0^\epsilon(X(0; x, t)), & \text{if } x > X(t; 0, 0); \\ u_1^\epsilon(S(t, x)), & \text{otherwise.} \end{cases}$$

We have the following result :

**Proposition 3.4**

The sequence  $\{u^n\}_{n \geq 0}$  converges strongly in  $L^1(\Omega_T)$  to the weak solution of problem (3.1) given by (3.23) .

**Proof.** We proceed as in [1], let  $\Omega_0^n, \Omega_1^n, \Omega_0, \Omega_1$  be the sets defined by

$$\begin{aligned} \Omega_0^n &= \{(t, x) \in \Omega_T; \quad x > X^n(t; 0, 0)\}, \\ \Omega_1^n &= \{(t, x) \in \Omega_T; \quad x < X^n(t; 0, 0)\}, \\ \Omega_0 &= \{(t, x) \in \Omega_T; \quad x > X(t; 0, 0)\}, \\ \Omega_1 &= \{(t, x) \in \Omega_T; \quad x < X(t; 0, 0)\}. \end{aligned}$$

and their respective characteristic functions  $\chi_0^n, \chi_1^n, \chi_0$  and  $\chi_1$ .

By virtue of (3.19), (3.21) we have :

$$\begin{aligned} u^n(t, x) - u(t, x) &= (u_0^\epsilon(X^n) - u_0^\epsilon(X))\chi_0^n \\ &\quad + (u_1^\epsilon(S^n) - u_1^\epsilon(S))\chi_1^n \\ &\quad + u_0^\epsilon(X)(\chi_0^n - \chi_0) \\ &\quad + u_1^\epsilon(S)(\chi_1^n - \chi_1) \end{aligned}$$

Note that

$$(3.24) \quad \chi_i^n \rightarrow \chi_i \quad \text{in} \quad L^1(\Omega_T) \quad i = 1, 2.$$

This follows from (3.18) and the fact that

$$\int_{\Omega_T} |\chi_i^n - \chi_i| dx dt \leq \int_0^T |\chi_i^n(t; 0, 0) - \chi_i(t; 0, 0)| dx dt.$$

It is clear that if  $u_0^\epsilon$  and  $u_1^\epsilon$  are regular functions then  $u^n \rightarrow u$  in  $L^2(\Omega_T)$ . Otherwise we proceed by regularization of the data  $u_0^\epsilon$  and  $u_1^\epsilon$  by considering two sequences  $u_{0,m}^\epsilon$  and  $u_{1,m}^\epsilon$  such that

$$(3.25) \quad \begin{cases} u_{0,m}^\epsilon \rightarrow u_0^\epsilon & \text{in } L^1(\Omega), \\ u_{1,m}^\epsilon \rightarrow u_1^\epsilon & \text{in } L^1(0, T) \end{cases}.$$

$$\begin{aligned} \int_{\Omega} |u_0^\epsilon(X^n) - u_0^\epsilon(X)| dx &\leq \int_{\Omega} |u_0^\epsilon(X^n) - u_{0,m}^\epsilon(X^n)| dx \\ &+ \int_{\Omega} |u_{0,m}^\epsilon(X^n) - u_{0,m}^\epsilon(X)| dx + \int_{\Omega} |u_{0,m}^\epsilon(X^n) - u_0^\epsilon(X)| dx \end{aligned}$$

The second term of the left hand side tends to zero as  $m \rightarrow +\infty$  by virtue of the regularity of  $u_0^\epsilon$  and (3.24).

To evaluate the first term, the change of variable

$$y = X^n(0; x, t), \quad x = X^n(t; y, 0)$$

leads to the inequality :

$$\int_{\Omega} |u_0^\epsilon(X^n) - u_{0,m}^\epsilon(X^n)| dx \leq \int_{\Omega} |u_0^\epsilon(y) - u_{0,m}^\epsilon(y)| \left| \frac{\partial}{\partial y} X^n(t; y, 0) \right| dy.$$

The a-priori estimate (3.14), Lemma 3.3 and (3.25) imply that

$$\int_{\Omega} |u_0^\epsilon(X^n) - u_{0,m}^\epsilon(X^n)| dx \rightarrow 0 \quad m \rightarrow +\infty,$$

and

$$\int_{\Omega} |u_0^\epsilon(X) - u_{0,m}^\epsilon(X)| dx \rightarrow 0 \quad m \rightarrow +\infty,$$

hence

$$\int_{\Omega} |u_0^\epsilon(X^n) - u_0^\epsilon(X)| dx \rightarrow 0 \quad n \rightarrow +\infty.$$

Following the same pattern and using the properties of  $S^n$  and  $S$  we deduce the convergence of  $u^n$  to  $u$  in  $L^1(\Omega_T)$ .

#### 4. The passage to the limit $\epsilon \searrow 0$ in problem $H_\epsilon$

The results of section 3 imply that the characteristic curves  $X^\epsilon$  and functions  $S^\epsilon$  defined by (3.21) lie in a bounded set of the space  $V_\delta$  for each  $\delta > 0$ . Thus two subsequences also denoted  $X^\epsilon, S^\epsilon$  converging in  $L^1(\delta, T; L^\infty(\Omega))$  can be extracted :

$$(4.1) \quad X^\epsilon \rightarrow X, \quad S^\epsilon \rightarrow S.$$

Assume that

$$(4.2) \quad u_i^\epsilon \rightarrow u_i \quad i = 0, 1$$

in a sense that we shall make precise in the sequel.

Let us define a function  $u$  by

$$(4.3) \quad u(t, x) = \begin{cases} u_0(X(0; x, t)), & \text{if } x > X(t; 0, 0); \\ u_1(S(t, x)), & \text{otherwise.} \end{cases}$$

Our objective is to establish the link between  $u^\epsilon$  and  $u$ .

First we have the

**Lemma 4.1**

The functions  $X$  and  $S$  given by (4.1) satisfy

$$(4.4) \quad \begin{cases} \Psi(X) = \int_s^t q(\sigma, X(\sigma; x, s)) d\sigma + \Psi(x) \\ X(S(t, x); x, t) = 0 \quad \text{a.e. in } \Omega_T \end{cases}$$

**Proof.** From properties (2.2) and (2.16), we have

$$\Psi^\epsilon(x) = \int_0^x \phi^\epsilon(\xi) d\xi \rightarrow \Psi(x) = \int_0^x \phi(\xi) d\xi \quad \text{a.e. } x \in \Omega$$

as  $|\Psi^\epsilon(x) - \Psi^\epsilon(y)| \leq \phi^+ |x - y|$  for  $x, y \in \Omega$ .

then (4.1) implies that

$$(4.5) \quad \Psi^\epsilon(X^\epsilon) \rightarrow \Psi(X) \quad \text{a.e. } (t, x) \in \Omega_T.$$

From (2.11) the property

$$(4.6) \quad |q^\epsilon(t, x) - q^\epsilon(t, y)| \leq a(t) |x - y| \quad \text{for } x, y \in \Omega \quad \text{holds.}$$

This allow us to pass to the limit in the term  $\int_s^t q^\epsilon(\sigma, X^\epsilon(\sigma; x, s)) d\sigma$ .

Indeed

$$\begin{aligned} R^\epsilon &= \int_s^t q^\epsilon(\sigma, X^\epsilon(\sigma; x, s)) - q(\sigma, X(\sigma; x, s)) d\sigma, \\ |R^\epsilon| &\leq \int_s^t |q^\epsilon(\sigma, X^\epsilon(\sigma; x, s)) - q^\epsilon(\sigma, X(\sigma; x, s))| d\sigma \\ &\quad + \int_s^t |q^\epsilon(\sigma, X(\sigma; x, s)) - q(\sigma, X(\sigma; x, s))| d\sigma \end{aligned}$$

as  $q^\epsilon(t, x) \rightarrow q(t, x)$  almost everywhere in  $\Omega_T$ .

As shown in (2.18) the second term tends to zero, the first one is bounded by

$$\int_s^t a(\sigma) |X^\epsilon(\sigma; x, s) - X(\sigma; x, s)| d\sigma$$

thus the convergence of  $R^\epsilon$  as  $\epsilon \searrow 0$  is assured and the validity of the first part of (4.4) is proved.

The second identity is deduced from (4.1) in a straightforward manner.

Now assume that

$$(4.7) \quad \begin{cases} u_0^\epsilon \rightarrow u_0 & \text{in } L^1(\Omega), \\ u_1^\epsilon \rightarrow u_1 & \text{in } L^1(0, T), \\ 0 \leq u_0, u_1 \leq 1. \end{cases}$$

Then the following convergence result is obtained

#### Theorem 4.2

The sequence  $\{u^\epsilon\}_{\epsilon>0}$  of solutions of problem (3.1) defined by (3.23) converges almost everywhere in  $\Omega_T$  to the function  $u$  given by (4.3).

**Proof.** Note that the sequence considered is that associated with the sequences  $X^\epsilon, S^\epsilon$  defined by (4.1).

Let  $I_\delta = \int_\delta^T \int_\Omega |u^\epsilon(t, x) - u(t, x)| dx dt$ .

First we have the decomposition

$$\begin{aligned} \chi_0^\epsilon u_0^\epsilon(X^\epsilon) - \chi_0 u_0(X) &= \chi_0^\epsilon u_0^\epsilon(X^\epsilon) - \chi_0 u_0^\epsilon(X^\epsilon) \\ &+ \chi_0 u_0^\epsilon(X^\epsilon) - \chi_0 u_0(X^\epsilon) + \chi_0 u_0(X^\epsilon) - \chi_0 u_0(X), \\ \chi_1^\epsilon u_1^\epsilon(S^\epsilon) - \chi_1 u_1(S) &= \chi_1^\epsilon u_1^\epsilon(S^\epsilon) - \chi_1 u_1^\epsilon(S^\epsilon) \\ &+ \chi_1 u_1^\epsilon(S^\epsilon) - \chi_1 u_1(S^\epsilon) + \chi_1 u_1(S^\epsilon) - \chi_1 u_1(S), \end{aligned}$$

so as in Lemma 3.3, Proposition 3.4 with properties (4.1) and (4.7) imply the different terms of  $I_\delta$  tend to zero in  $L^1(\delta, T; L^1(\Omega))$  when  $\epsilon \searrow 0$ .

In the more general case we are looking for equations satisfied by the limit as  $\epsilon \searrow 0$  of the sequences  $\{u^\epsilon\}_{\epsilon>0}$ .

The sequence  $\{u^\epsilon\}_{\epsilon>0}$  defined by (3.23) is uniformly bounded in  $L^\infty(\Omega_T)$ , so we can substract a sequence that converges to  $u$  in  $L^\infty(\Omega_T)$  weak\*.

We assume that the hypothesis (2.12) holds. We have

$$(4.8) \quad \begin{cases} \frac{\partial}{\partial t}(\phi^\epsilon p^\epsilon) + \frac{\partial}{\partial x} q^\epsilon = 0, \\ \frac{\partial}{\partial t}(\phi^\epsilon u^\epsilon) + \frac{\partial}{\partial x}(q^\epsilon u^\epsilon) + \phi^\epsilon u^\epsilon \frac{\partial}{\partial t} p^\epsilon = 0. \end{cases}$$

From (2.2), (2.15) and (3.23) it follows that  $F^\epsilon = \phi^\epsilon u^\epsilon \frac{\partial}{\partial t} p^\epsilon$  belongs to a bounded set of  $L^2(\Omega_T)$ . If we denote by

$$A^\epsilon = (\phi^\epsilon u^\epsilon, q^\epsilon u^\epsilon), \quad B^\epsilon = (-q^\epsilon, \phi^\epsilon p^\epsilon)$$

we have

$$\operatorname{curl} B^\epsilon = F^\epsilon, \quad \operatorname{div} A^\epsilon = 0$$

Using the elegant method of compensated compactness initiated by MURAT [20] and TARTAR [28] we have that

$\langle A^\epsilon, B^\epsilon \rangle$  converges to  $\langle A, B \rangle$  in the sense of distributions.

The following assertions are valid

$$\langle A^\epsilon, B^\epsilon \rangle = (q^\epsilon u^\epsilon)(\phi^\epsilon p^\epsilon) - (q^\epsilon)(\phi^\epsilon u^\epsilon),$$

$$q^\epsilon \rightarrow q \quad \text{a.e. in } \Omega_T,$$

$$p^\epsilon \rightarrow p \quad \text{a.e. in } \Omega_T,$$

$$(\phi^\epsilon u^\epsilon) \rightharpoonup \widetilde{\phi u} \quad \text{in } L^\infty(\Omega_T) \quad \text{weak*},$$

$$(\phi^\epsilon q^\epsilon) \rightharpoonup \phi q \quad \text{in } L^\infty(\Omega_T) \quad \text{weak*},$$

$$u^\epsilon \rightharpoonup u \quad \text{in } L^\infty(\Omega_T) \quad \text{weak*}$$

which implies

$$\langle A^\epsilon, B^\epsilon \rangle \rightharpoonup \widetilde{\phi u} q - \widetilde{q u} \widetilde{\phi p} \quad \text{in the distributional sense.}$$

We have also

$$\langle A^\epsilon, B^\epsilon \rangle \rightharpoonup (\phi q) u - \widetilde{q u} \widetilde{\phi p},$$

where the uniqueness of the limit in  $\mathcal{D}'(\Omega_T)$  implies that

$$\widetilde{\phi u} = \phi u.$$

Considering (2.16), we have

$$\frac{\partial}{\partial t} p^\epsilon \rightarrow \frac{\partial}{\partial t} p \quad \text{a.e. in } \Omega_T,$$

where  $p$  is the solution of the homogenized problem (2.1) related to  $q$  by

$$\phi(x) \frac{\partial}{\partial t} p + \frac{\partial}{\partial x} q = 0, \quad q = -k \frac{\partial}{\partial x} p$$

$$F^\epsilon \rightharpoonup \phi u \frac{\partial}{\partial t} p \quad \text{in } L^2(\Omega_T) \quad \text{weak}.$$

In a distributional sense  $u$  satisfies

$$\phi(x) \frac{\partial}{\partial t} p + q(t, x) \frac{\partial}{\partial x} u = 0 \quad \text{in } \Omega_T,$$

which completes the proof.

## 5. References

- [1] Y. AMIRAT, K. HAMDACHE and A. ZIANI,  
Homogénéisation d'un modèle d'écoulements miscibles en milieu poreux,  
Rapport INRIA n° 802 (1988).
- [2] Y. AMIRAT, K. HAMDACHE and A. ZIANI,  
Homogénéisation d'équations hyperboliques du premier ordre;  
Application aux milieux poreux Rapport INRIA n° 803 (1988).
- [3] J.L. AURIAULT and E. SANCHEZ PALENCIA, Remarques sur la loi de Darcy pour  
les écoulements biphasiques en milieu poreux, *Journal de Mécanique théorique et  
Appliquée* (1986) (special issue), 141-156.
- [4] N.S. BAKHVALOV and G.P. PANASENKO,  
*Processus de moyenne dans les milieux periodiques* (Nauka, Moscow, 1984) (in Russian)
- [5] J. BEAR, *Dynamics of fluids in porous media*, (Elsevier, 1972).
- [6] A. BENSOUSSAN, J.L. LIONS and G.C. PAPANICOLAOU,  
*Asymptotic Analysis for periodic structures*, (North-Holland, 1978).
- [7] A. BOURGEAT, Homogenized behavior of two phase flows in naturally fractured  
reservoirs with uniform fractures distributions, *Computer Methods in Applied Me-  
chanics and Engineering* 47 (1984) 205-216.
- [8] R.E. CAFLISCH and J. RUBINSTEIN, Lecture on mathematical theory of multiphase  
flow, (Courant Inst. of Math. Sciences, 1984)
- [9] G. CHAVENT and J. JAFFRE,  
*Mathematical Models and Finite Elements for Reservoir Simulation*,  
(North-Holland, 1986)
- [10] H. DARCY, *Les fontaines publiques de la ville de Dijon* (Dalmon, Paris, 1856).
- [11] J. DOUGLAS and J.E. ROBERTS, Numerical Methods for a model for compressible  
miscible displacement in porous media, *Math. of Computation* 41, n° 164, 1984.
- [12] R. FIGARI, G.C. PAPANICOLAOU and J. RUBINSTEIN, The point interaction ap-  
proximation for diffusion in regions with small holes, (Reprint)
- [13] A. HOUPERT,  
*Mécanique des fluides dans les milieux poreux : Critiques et Recherches*,  
(Technip, Paris, 1972).
- [14] O.A. LADYZENSKAJA, V.A. SOLONNIKOV and N.N. URAL'CEVA,  
*Linear and Quasilinear Equations of Parabolic Type*  
(Trans. of Math. Monographs R.I. 28, 1968).
- [15] J.L. LIONS  
*Some methods in the mathematical analysis of systems and their control*,  
(Gordon and Beach, 1981).
- [16] V.A. MARCHENKO and E.J. HROUSLOV,  
*Boundary Value Problems in domains with granuled boundaries*,  
(Nauka-Dumka, Kiev, 1974).
- [17] V.G. MARKOV and O.A. OLEINIK, On propagation of heat in one dimensional  
disperse media, *Journal of Appl. Math. Mech.*, 37 (1975) 1028-1037.

- [18] C.M. MARLES,  
*Les écoulements polyphasiques en milieu poreux*, (Masson, Paris, 1967).
- [18] G. MATHERON, *Eléments pour une théorie des milieux poreux* (Masson, Paris, 1967).
- [20] F. MURAT,  
a- H-convergence, Séminaire Facultés des Sciences, Alger, (1978).  
b- Compacité par compensation, Ann. Scuola Norm. Sup. Pisa, V. 5, 1978, 489-507.
- [21] M. MUSKAT,  
a- *The flow of Homogeneous Fluids through porous media* (Mc Graw Hill N.Y. 1937).  
b- *Principles of oil production* (Mc Graw Hill N.Y., 1949).
- [22] J. RUBINSTEIN, On the macroscopic description of slow viscous flow past a random array of spheres, (Reprint).
- [23] E. SANCHEZ PALENCIA,  
a- *Non Homogeneous media and Vibration theory*,  
(Lectures Notes in Physics, 127, 1980).  
b- Homogenization in Mechanics : A survey of solved and open problems, Rend. sem. mat. 44-1 Torino (1986).
- [24] A.E. SCHEIDEGGER, *The Physics of Flow through porous media* (Univ Toronto, 1974).
- [25] M.I. SHVIDLER,  
a- Dispersion of a filtration stream in a medium with random inhomogeneities, Soviet Phys. Dokl. 20 n° 3, (1975).  
b- Averaging Transfer Equations in porous media with random inhomogeneities, Transl from Izv. Akad Nauk S.S.S.R., Mekh. Zhidk., Gaza I. 1 (1985).
- [26] J.C. SLATTERY,  
*Momentum, Energy and Mass Transfer in Continua* (Mc Graw Hill N.Y., 1972).
- [27] S. SPAGNOLO, Convergence of Parabolic Equations, Bolletino U.M.I.(5), 14-B (1977).
- [28] L. TARTAR,  
a- Cours Peccot, Collège de France (1977).  
b- Compensated Compactness and Applications to P.D.E., in *Research Notes in Math., Non Linear Analysis and Mechanics*, Heriot-Watt Symposium, 4, V. 39, Ed R.J. KNOPS, Pitman Press, 1979.  
c- Non-local effects induced by Homogenization, in *Essays of mathematical analysis in honour of E. De GIORGI* (Birkhauser 1989).



